

CHRONIC OBSTRUCTIVE PULMONARY DISEASE



Chronic coughing?

Coughing more than usual



Out of breath?

Frequent shortness of breath



Passive smoke exposure?

Exposure to secondhand smoke



Daily smoking habits?

Constant and regular smoking

ASSESS YOUR RISK FOR COPD.

Contact your GP or your Community
Health Service now for a consult.

REFERENCES:

1. Respiratory Disease in New Zealand. Available from <https://www.asthmafoundation.org.nz/research/key-statistics> (Accessed May 2019)
2. Young, Robert. "Management of chronic obstructive pulmonary disease (COPD) – with a focus on the 2017 GOLD Strategy update." A Research Review™ Educational Series, 2017, p.2
3. Young, Robert. "Management of chronic obstructive pulmonary disease (COPD) – with a focus on the 2017 GOLD Strategy update." A Research Review™ Educational Series, 2017, p.2
4. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease, 2019 Report. Pages 53 and 92. Accessed on April 2019.

Duolin Inhaler, PRESCRIPTION MEDICINE. Duolin Inhaler contains Salbutamol (as sulphate) 100 mcg and Ipratropium bromide 20 mcg per actuation. Each Duolin Inhaler contains 200 actuations. Duolin Inhaler is used for the treatment of obstructive airway diseases such as COPD, bronchitis and emphysema.

Duolin Inhaler has risks and benefits. Do not use Duolin Inhaler if you are allergic to salbutamol, ipratropium bromide, or any of the other ingredients of Duolin Inhaler. In patients who find coordination of a pressurised metered dose inhaler difficult, it may be better for you to use a spacer device with your Duolin Inhaler. Ask your doctor or your pharmacist to explain how to use it, or follow the instructions provided with the spacer.

Always use Duolin Inhaler exactly as your doctor or respiratory nurse has told you. Duolin Inhaler is for oral inhalation only. If you have any concerns, discuss them with your doctor or respiratory nurse. Discuss your asthma or chronic obstructive pulmonary disease (COPD) symptoms with your doctor or pharmacist and ask whether Duolin Inhaler is the appropriate choice for you.

Duolin Inhaler is fully funded. Standard doctor's fees and pharmacy charges apply.

Duolin Inhaler is distributed by REX Medical Ltd, Auckland, New Zealand.

© REX Medical Ltd

TAPS NA 11116 | CC190531-V1



IN NEW ZEALAND,
COPD AFFECTS AN ESTIMATED

1 IN 7
ADULTS
aged over 45 years¹



FOR EVERY
4 - 5 PATIENTS
WITH COPD,
ONLY 1
WILL HAVE
A CONFIRMED
DIAGNOSIS²



**COPD LUNG DAMAGE
IS IRREVERSIBLE**

By the time a diagnosis is
confirmed, approximately

50% OF
LUNG FUNCTION
may be irrevocably
damaged³

CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD)

C

Coughing more than usual?

O

Out of breath frequently?

P

Passive exposure to
secondhand smoke?

D

Daily smoking habits?

ASSESS YOUR RISK FOR COPD.

Contact your GP or your Community
Health Service now for a consult.

Duolin Inhaler (Salbutamol / Ipratropium) is appropriate for regular and as-needed use to improve symptoms in patients with mild COPD requiring no hospitalisation. When it comes to improving FEV₁ and symptoms, the combination of SABA (Salbutamol) and SAMA (Ipratropium) is found to be superior than either medication alone.⁴

Consult with your GP now and ask about **Duolin Inhaler**,
your first defence against COPD.

Duolin Inhaler is indicated for the treatment of obstructive airway diseases such as COPD, bronchitis and emphysema.

